

Purpose: Ran 6% Tris - Glycine gel of *Brachamella catarrhalis* 49143 OG extract (KT-frozen cells - 11/14/94) and silver stained to determine approximate size of OMP106 protein or homolog.

Methods:

1. Cast continuous 6% Tris - Glycine gel; SDS-PAGE.
2. Prepared samples:
 - Bio-Rad Pre-stained (Low Range)
 - Bio-Rad Silver Standard (Broad Range) 1:20 dilutions
 - Novex standard
 - OG - B. cat. 49143 (Sample labeled KT-frozen cells - 11/14/94) 1:2 dilution w/ 2X SB; one sample was boiled, other sample was not boiled
3. Ran gel @ 20 mAmp constant current.
4. Silver stained gel.

Results:

Lane - sample

- 1 - BioRad Pre-stained Low Range
- 2 - OG Not Boiled (25ul of 1:2 diluted sample)
- 3 - OG Boiled (25ul of 1:2 diluted sample)
- 4 - Novex Std.
- 5 - OG Not Boiled
- 6 - BioRad Silver Standard Broad Range (25ul of 1:20 diluted sample)
- 7 - OG Boiled
- 8 - BioRad Pre-stained Low Range

Standards MW (Daltons):

Bio-Rad Pre-stained (Low)

139,900
86,800
47,800
33,300
28,600
20,100

Novex

250,000
98,000
64,000
50,000
36,000

Bio-Rad Silver Std (Broad)

200,000
116,250
97,400
66,200

Continued on Page

Read and Understood By

Holly Dilbert

Signed

/ Date

Signed

Date

Conclusion:

The size of the OMP106 is approx. 200 kDa. The protein does not migrate into gel unless boiled; compare boiled and not boiled lanes for protein of interest.

Continued on Page

Read and Understood By

Holly Obeir

Signed

Date

J. Obeir

Signed

Date



RECEIVED
OCT 17 2003
TECH CENTER 1600/2900
BioRad STD



Unboiled
extract

Boiled extract

Novex STD

Unboiled
extract

BioRad Sil.
STD

Boiled extract

BioRad STD